



Childhood exposure to constricted living space: a possible environmental threat for myopia development

Kai Yip Choi¹, Wing Yan Yu¹, Christie Hang I Lam¹, Zhe Chuang Li¹, Man Pan Chin¹, Yamunadevi Lakshmanan¹, Francisca Siu Yin Wong¹, Chi Wai Do¹, Paul Hong Lee²  and Henry Ho Lung Chan¹

¹The Centre for Myopia Research, School of Optometry, The Hong Kong Polytechnic University, Kowloon, Hong Kong, and ²School of Nursing, The Hong Kong Polytechnic University, Kowloon, Hong Kong

Citation information: Choi KY, Yu WY, Lam CHI, Li ZC, Chin MP, Lakshmanan Y, Wong FSY, Do CW, Lee PH & Chan HHL. Childhood exposure to constricted living space: a possible environmental threat for myopia development. *Ophthalmic Physiol Opt* 2017; 37: 568–575. <https://doi.org/10.1111/opo.12397>

Keywords: childhood refractive error, constricted living space, epidemiology, living environment, myopia, myopia prevalence

Correspondence: Henry H L Chan
E-mail address: henryhl.chan@polyu.edu.hk

Received: 28 November 2016; Accepted: 16 May 2017; Published Online: 22 June 2017

Abstract

Purpose: People in Hong Kong generally live in a densely populated area and their homes are smaller compared with most other cities worldwide. Interestingly, East Asian cities with high population densities seem to have higher myopia prevalence, but the association between them has not been established. This study investigated whether the crowded habitat in Hong Kong is associated with refractive error among children.

Methods: In total, 1075 subjects [Mean age (S.D.): 9.95 years (0.97), 586 boys] were recruited. Information such as demographics, living environment, parental education and ocular status were collected using parental questionnaires. The ocular axial length and refractive status of all subjects were measured by qualified personnel.

Results: Ocular axial length was found to be significantly longer among those living in districts with a higher population density ($F_{2,1072} = 6.15, p = 0.002$) and those living in a smaller home ($F_{2,1072} = 3.16, p = 0.04$). Axial lengths were the same among different types of housing ($F_{3,1071} = 1.24, p = 0.29$). Non-cycloplegic autorefractometry suggested a more negative refractive error in those living in districts with a higher population density ($F_{2,1072} = 7.88, p < 0.001$) and those living in a smaller home ($F_{2,1072} = 4.25, p = 0.02$). After adjustment for other confounding covariates, the population density and home size also significantly predicted axial length and non-cycloplegic refractive error in the multiple linear regression model, while axial length and refractive error had no relationship with types of housing.

Conclusions: Axial length in children and childhood refractive error were associated with high population density and small home size. A constricted living space may be an environmental threat for myopia development in children.

Introduction

Myopia, or short sightedness, which is the most common refractive error, can be regarded as a type of ocular disorder. It has been a global health concern with costs including not only optical corrections to obtain clear distant vision, but also the medical burden of high myopes who are pre-disposed to various ocular diseases such as cataract, glaucoma, macular degeneration and retinal detachment¹

which can cause severe or irreversible vision loss. As vision is crucial to our daily life, the vision loss can adversely affect the quality of life.²

‘Emmetropization’ is a visually guided process for the eye to modify itself to obtain an optimum relationship between the axial length and other ocular components, such as the cornea and lens, so that any infantile refractive error is corrected. However, there is an increasing number of children becoming myopic at an early age. Not only will

this increase their risk of developing high myopia later in their lives, but epidemic childhood myopia is also speculated to cause a shortage of certain labour forces as good uncorrected eyesight is a pre-requisite for some occupations such as pilots and firefighters, and thus lead to an increasing social burden in the coming decades.

East Asian countries generally have an unexpectedly higher myopia prevalence compared with other parts of the world.³ Among them, Hong Kong has long been a city with an extraordinary high prevalence of myopia.^{4,5} Studies have shown that myopia is more prevalent in Asians than in white European and African populations.^{6,7} Apart from genetic differences, these findings were found to be associated with the culture and lifestyle of East Asians, who are usually lacking in outdoor activities and engaged in a near-work-predominant education system.^{6,8} In addition, the crowded living habitat among the East Asian cities may also be associated with this high prevalence of myopia.

Previous studies revealed that an urban environment is related to a higher prevalence of myopia in children compared with sub-urban and rural environments.^{9,10} For example, the Sydney Myopia Study (SMS) suggested that the urbanicity of the living region was associated with childhood myopia,¹¹ in which the children living in a place of denser population were reported to have a higher prevalence of myopia. Some other studies attributed the association to the lack of outdoor activities and the excess of near work^{12–14} for children living in an urban area. The Sydney Myopia Study also reported that flat-styled rather than house-styled living in an urban area had an association with myopia prevalence. A recent study also suggested that the taller the building that the children were living in, the higher the chance that myopia would be observed.¹⁵

In 2004, Fan *et al.* conducted a population-based study on myopia prevalence in Hong Kong, which included 7560 schoolchildren.⁴ From the results, 37% of the children were found to be myopic. They recruited one school from each of the 18 political districts in Hong Kong. However, among the 18 political districts, half of them had a population density lower than 10 000 persons per km², while only a few of them had a population density higher than 30 000 persons per km².¹⁶ Their samples may have been skewed towards the less populated areas and thus they may have underestimated the actual myopia prevalence of Hong Kong according to SMS.¹¹

Hong Kong is one of the most densely populated cities in East Asia. The housing problem in Hong Kong has been intensely discussed, as the land supply is limited while the population is increasing.¹⁶ In 2015, 45.7% of the Hong Kong population lived in public housing,¹⁷ and the internal floor area per person was only 13.1 m². While there are still hundreds of thousands of people queuing for public housing, it was reported that around 171 000 people in Hong

Kong live in substandard sub-divided flats.¹⁸ Some children even have to live in flats with a total area of around 9 m² with their whole family.¹⁸

In East Asian cities, people are generally living in relatively small flats in highly populated areas and the prevalence of myopia is high. However, the association between refractive error and size of living space has not been established. In the current study, we studied whether this crowded living environment is associated with refractive error among children in primary (elementary) school.

Methods

Subjects

Local primary school children were recruited between June 2015 and February 2016. Cluster sampling was used for the selection of schools. The 18 political districts in Hong Kong were divided into three clusters according to their population densities¹⁶: high: more than 30 000 persons per km²; medium: 10 000 to 30 000 persons per km²; and low: less than 10 000 persons per km². In each cluster, four schools were randomly selected (12 schools in total). Eight schools finally agreed to join the study (two schools from the low density cluster, three schools from the medium density cluster, and three schools from the high density cluster). All students who fulfilled the inclusion criteria were examined in a vision screening which was held in their school. Inclusion criteria were students aged from 7 to 12 years who were a Hong Kong Chinese resident. In total, 1235 students were invited to join the study, and 1173 students participated (95% response rate). Among them, 19 students exceeded the age limit and 15 mainland China residents were excluded from the study. Furthermore, we excluded 64 respondents from the analysis who had received different active myopia control interventions. As a result, 1075 [Age (S.D.): 9.95 (0.97) years, 54.5% boys] students were included in the current study for analysis. As all the subjects were local Chinese residents studying in government-supported schools with the same syllabus governed by the Education Bureau of The Hong Kong Government, we assumed that all subjects received a similar education which would not differ between groups. Informed consent and simple written assent were obtained from the parents and the students respectively. All procedures followed the tenet of Declaration of Helsinki and were approved by the Human Subjects Ethics Subcommittee of The Hong Kong Polytechnic University.

Data collection

The demographic data, ocular and family history, parental education level, and information regarding living environment were obtained by a self-reported questionnaire, which

was completed by the parents. For the living environment, information of the residential district, the home size and the physical type of housing were collected. Different home sizes were categorised as smaller than 27.87 m² (300 ft²), from 27.87 to 55.74 m² (from 300 to 600 ft²) and larger than 55.74 m² (600 ft²), which were based on the common living style in Hong Kong. For the residential district, we grouped them according to their population densities¹⁶ into low, medium and high population density which were defined as less than 10 000, from 10 000 to 30 000 and more than 30 000 persons per km² in the district respectively.

Ocular axial length (AL, length of eyeball) of the subjects was measured using partial coherence interferometry (Carl Zeiss Meditec, IOL Master, <https://www.zeiss.com/meditec/int/products/ophthalmology-optometry/cataract/diagnostic/s/optical-biometry/iolmaster-500.html>). A total of five measurements (signal-to-noise ratio > 2.0) were taken and the mean value was recorded. Their refractive status was evaluated by non-cycloplegic open-field auto-refraction (Shin-Nippon, NVision K5001, <http://www.shin-nippon.jp/products/nvk5001/>) while looking at a distant target at 6 m. Cycloplegic agents were not instilled because the data were collected on normal school days, and we did not want to interrupt the students' daily study. This is one of the limitations of our study as the students may accommodate, resulting in a myopia over-estimation in the auto-refraction results.¹⁹ A total of five measurements within the published criteria were taken by an optometrist, and the representative value was recorded.²⁰ The recorded value was then transposed into spherical equivalent refraction (SER) by the following equation: SER = spherical refractive error + " cylindrical refractive error.

Statistical analysis

Data analysis was performed using SPSS (IBM, ver. 22, <https://www.ibm.com/analytics/us/en/technology/spss/>).

Axial length was the primary outcome and non-cycloplegic spherical equivalent refraction was the secondary outcome to assess the characteristics and trends between groups. Each independent variable was plotted against AL and SER, and the results were compared among groups using one-way ANOVA with Bonferroni correction. Multiple linear regression was used to assess the impact of population density, home size, and type of housing on AL and SER. Confounding covariates included age, gender, parental education level, and parental myopia. Missing data were treated using 10-time multiple imputation.²¹ To ensure the absence of multicollinearity, only models showing the following signs (all collinearity tolerances larger than 0.8, all variance inflation factors less than 2 and all absolute Pearson's *R* of variables was smaller than 0.2) were analysed. As

data from right and left eyes were strongly correlated (AL: $r = 0.96$; SER: $r = 0.92$), only right eye data were analysed. Significance level was set as $p < 0.05$.

Results

Descriptive characteristics of the sample

The subjects had a mean AL of 23.78 mm (S.D.: 1.04) and SER of -1.21 D (S.D.: 1.80). *Table 1* shows the demographics and living environment of the participants, and the p values in *Table 1* were from a univariate analysis of each variable. The age of the children did not significantly differ across all categories of population density ($F_{2,1072} = 2.82$, $p = 0.06$), home size ($F_{2,1072} = 2.10$, $p = 0.12$), and type of housing ($F_{3,1071} = 1.60$, $p = 0.19$).

Living environment – between group comparison

AL and SER were plotted across different groups of each variable individually. For AL, we observed significant difference in population density of the residential district ($F_{2,1072} = 6.15$, $p = 0.002$, *Figure 1*) and home size ($F_{2,1072} = 3.16$, $p = 0.04$, *Figure 2*). However, the difference in association of AL in type of housing was not significant ($F_{3,1071} = 1.24$, $p = 0.29$). Axial length increased as population density of the residential districts increased, but significant difference could only be observed in districts with low population density when compared with those with high population density ($p = 0.002$). There was also a decreasing trend of AL with home size. A significant difference was observed between those living in a larger home and those living in a smaller home ($p = 0.04$). For SER, we also observed significant difference in population density of the residential district ($F_{2,1072} = 7.88$, $p < 0.001$, *Figure 1*) and home size ($F_{2,1072} = 4.25$, $p = 0.02$, *Figure 2*). However, the difference in association of SER in type of housing was again insignificant ($F_{3,1071} = 1.75$, $p = 0.16$). SER was more negative as population density of the residential districts increased. Significant difference could be observed in districts with low population density when compared with those with high population density ($p = 0.001$) and districts with medium population density when compared with those with high population density ($p = 0.009$). SER was less negative as home size increased. A significant difference was observed between those living in a large-sized home and those living in a small-sized home ($p = 0.02$), and between those living in a large-sized home and those living in a medium-sized home ($p = 0.03$).

Living environment – multivariate analysis

The multiple linear regression models were overall significant (AL: $F_{14,1060} = 10.26$, $p < 0.001$; SER: $F_{14,1060} = 4.88$,

Table 1. Distribution of demographics and living environment factors

	N (%)	Mean AL (S.D.)	p value^a	Mean SER (S.D.)	p value^a
All	1075 (100)	23.78 (1.04)		-1.21 (1.80)	
Gender					
Boys	586 (54.5)	24.02 (1.00)	<0.001	-1.20 (1.80)	0.94
Girls	489 (45.5)	23.49 (1.02)		-1.21 (1.80)	
Age					
Lower third	358 (33.3)	23.53 (0.93)	<0.001	-0.90 (1.64)	<0.001
Middle third	358 (33.3)	23.80 (1.06)		-1.34 (1.85)	
Upper third	359 (33.4)	24.02 (1.07)		-1.41 (1.87)	
Parental myopia					
No parent is myopic	507 (47.2)	23.71 (1.02)	<0.001	-1.00 (1.59)	<0.001
One parent is myopic	336 (31.3)	23.83 (1.07)		-1.37 (1.94)	
Both parents are myopic	152 (14.1)	24.09 (1.11)		-1.86 (2.01)	
Parental education level					
Primary school or below	58 (5.4)	24.06 (1.07)	0.11	-1.53 (2.12)	0.51
Junior secondary school	375 (34.9)	23.78 (1.05)		-1.22 (1.76)	
Senior secondary school	422 (39.3)	23.72 (1.02)		-1.17 (1.80)	
Tertiary education	163 (15.2)	23.83 (1.08)		-1.12 (1.65)	
Population density of the residential district					
<10k persons per km ²	209 (19.4)	23.56 (0.93)	0.002	-0.89 (1.64)	<0.001
10k–30k persons per km ²	236 (22.0)	23.74 (1.07)		-1.01 (1.60)	
>30k persons per km ²	418 (38.9)	23.87 (1.09)		-1.46 (2.01)	
Home size					
<27.87 m ² (<300 ft ²)	305 (28.4)	23.85 (1.07)	0.043	-1.35 (1.88)	0.015
27.87–55.74 m ² (300–600 ft ²)	536 (49.9)	23.80 (1.10)		-1.26 (1.89)	
>55.74 m ² (>600 ft ²)	152 (14.1)	23.59 (0.88)		-0.82 (1.38)	
Type of housing					
Flat	913 (84.9)	23.77 (1.05)	0.29	-1.22 (1.81)	0.16
Suite	38 (3.5)	24.00 (1.10)		-1.54 (1.83)	
House/Penthouse	29 (2.4)	23.52 (0.94)		-0.50 (1.65)	
Rooftop shack/Sub-divided unit	22 (2.0)	23.92 (1.27)		-1.16 (1.69)	

AL, Axial length; SER, Spherical equivalent refraction.

^a*p* values reported here were the significance level of univariate analysis between groups.

$p < 0.001$) and the adjusted R^2 were 0.13 and 0.06 respectively. Table 2 summarises the effect of individual variable, and the p values were from a multivariate analysis of all variables, adjusted for gender, age, parental education level and parental myopia. Among individual target covariates in the AL model, only population density of the residential district and home size made significant contributions. The B value (regression coefficient) for living in a district of high population density was 0.24 (95% CI 0.07–0.40), indicating that children living in districts with high population density were predicted to have a 0.24 mm longer eye compared with those living in districts with low population density. However, the B value for medium population density was not significant ($p = 0.45$). The home size recorded a B value of 0.25 (95% CI 0.05–0.46) when comparing a large-sized and a small-sized home, predicting a 0.25 mm longer eye for those living in a small-sized home. For a medium-sized home, the B value was 0.19 (95% CI 0.00–0.37) when compared to a large-sized home, indicating a

0.19 mm longer eye for those living in a medium-sized home. Furthermore, type of housing did not significantly contribute to the model (Suite: $p = 0.17$; House/Penthouse: $p = 0.95$; Sub-divided unit/Rooftop shack: $p = 0.94$). In the SER model, only population density of the residential district and home size showed significant contributions. The B value for living in a district of high population density was -0.53 (95% CI -0.83 to -0.23), indicating that the SER of children living in districts with high population density were predicted to be 0.53 D more myopic, or less hyperopic, compared with those living in districts with low population density. However, the B value for medium population density was not significant ($p = 0.90$). The home size recorded a B value of -0.47 (95% CI -0.86 to -0.08) when comparing a large-sized and a small-sized home, predicting the SER for those living in a small-sized home were 0.47 D less, but the B value for medium-sized home was not significant ($p = 0.06$). Furthermore, type of housing did not significantly contribute to the model (Suite:




Figure 1. Association of population density of the residential district with axial length (AL) and non-cycloplegic spherical equivalent refraction (SER). The triangles and squares represent the mean ± S.E. of AL and SER, respectively. Bonferroni correction: * $p < 0.05$, ** $p < 0.01$.




Figure 2. Association of home size with axial length (AL) and non-cycloplegic spherical equivalent refraction (SER). The triangles and squares represent the mean ± S.E. of AL and SER, respectively. Bonferroni correction: * $p < 0.05$, ** $p < 0.01$.

$p = 0.26$; House/Penthouse: $p = 0.63$; Sub-divided unit/Rooftop shack: $p = 0.72$). The non-cycloplegic SER was similar to and supported the AL results.

Discussion

The results of this study provide further support for an association between living environment and childhood refractive error. One of our major findings is that children living in districts of higher population density have a higher risk of having a longer eye and a more negative non-cycloplegic SER. Other research studies have also shown supporting results.^{9–11} The Refractive Error Study in Children (RESC)²² provided a standardised protocol to measure the prevalence of refractive error in school-aged children

worldwide,^{9,23–26} enabling easy comparison as all the sampling and measurement protocols were the same. The RESC group found that studies conducted in urban areas revealed a higher myopia prevalence than those in rural areas.^{9,27,28} Besides RESC, the Sydney Myopia Study²⁹ investigated many modifiable risk factors such as volume of near work,¹⁴ time spent in outdoor activities,³⁰ and urbanicity of the residence.¹¹ For the living environment, Ip and co-workers found that children living in the inner city were more likely to have myopia than those living in outer suburban areas. In Hong Kong, the results were similar. We grouped the 18 political districts in Hong Kong into three clusters according to their population densities¹⁶ and observed that population density was associated with the risk of having a longer eye (Figure 1). Similar trends were

Table 2. Multivariate analysis on axial length and spherical equivalent refraction

	B value (S.E.)	95% CI	p value
Axial length (mm)			
Population density of the residential district			
>30k persons per km ²	0.24 (0.08)	0.07 to 0.40	0.005
10k–30k persons per km ²	0.07 (0.09)	–0.11 to 0.25	0.45
(ref = <10k persons per km ²)			
Home size			
<27.78 m ²	0.25 (0.10)	0.05 to 0.46	0.01
27.78–55.74 m ²	0.19 (0.09)	0.00 to 0.37	0.05
(ref = >55.74 m ²)			
Type of housing			
Suite	0.22 (0.17)	–0.10 to 0.55	0.17
House/Penthouse	–0.01 (0.18)	–0.37 to 0.35	0.95
Rooftop shack/ Sub-divided unit	–0.02 (0.21)	–0.43 to 0.40	0.94
(ref = Flat)			
Spherical equivalent refraction (D)			
Population density of the residential district			
>30k persons per km ²	–0.53 (0.15)	–0.83 to –0.23	0.001
10k–30k persons per km ²	–0.02 (0.16)	–0.33 to 0.29	0.90
(ref = <10k persons per km ²)			
Home size			
<27.78 m ²	–0.47 (0.20)	–0.86 to –0.08	0.02
27.78–55.74 m ²	–0.31 (0.17)	–0.64 to 0.02	0.06
(ref = >55.74 m ²)			
Type of housing			
Suite	–0.35 (0.31)	–0.95 to 0.25	0.26
House/Penthouse	0.17 (0.34)	–0.51 to 0.84	0.63
Rooftop shack/ Sub-divided unit	0.14 (0.40)	–0.63 to 0.92	0.72
(ref = Flat)			

Confounding covariates included age, gender, parental myopia, and parental education level.

observed in both big and small cities, Sydney and Hong Kong, thus the effect of urbanicity ought not to be overlooked in considering factors that associate with childhood refractive error.

The second major observation of our study was the association of the home size with childhood refractive error. Children living in a home smaller than 27.87 m² (300 ft²) had a significantly longer eye when compared to those living in a home larger than 55.74 m² (600 ft²). Although myopia prevalence was thought to increase with socioeconomic status, which can partially be reflected by large home size and high parental education level, in our sample the small home size showed a stronger association with longer axial length and more negative SER than higher parental education level (AL: $F_{3,1071} = 2.02$, $p = 0.11$; SER: $F_{3,1071} = 0.77$, $p = 0.51$). One possible reason may be the

constricted environment at home creating peripheral hyperopic defocus from the surroundings. Numerous studies had shown that peripheral hyperopic defocus accelerates, while peripheral myopic defocus retards, myopia progression.^{31–34} In different visual environments, objects nearby produce various amount of defocus to the eye with regards to the plane of focus.^{35,36} Generally, an indoor environment creates more peripheral hyperopic defocus than an outdoor environment.³⁶ This condition may also apply to a constricted area in an indoor setting vs an open area, thus children in a smaller home would be exposed to stronger peripheral hyperopic defocus compared with those in a larger home.

The type of housing may be another factor associated with myopia prevalence. A recent nationwide population-based study in China evaluated the impact of living environment on myopia in school-aged children.¹⁵ From their sizable sample, myopia was associated with the type of housing, in terms of the height of residential buildings. Higher myopia prevalence was observed in children living in taller buildings, which is independent of the residential region, age, gender and ethnicity. In the Sydney Myopia Study, myopia was more frequently observed in children living in apartments and terrace houses than those living in stand-alone or separate houses.¹¹ They suggested it was related to the nature of housing type, among which terrace houses and apartments are smaller and more confined. However, studies in Singapore did not show such a relationship.^{37,38} Our study showed that home size was associated with axial length and refractive error instead of the type of housing. One possible reason for the insignificance may be the variation of housing type in Hong Kong was relatively too little, as the majority live in a flat-styled home. This could be a possible explanation why Asian children living in urban area are more likely to have myopia as they mostly live in flat-styled accommodation, yet this could not be determined in our study.

The housing issue has been a complicated problem in Hong Kong. In 2015, the average living space per person in public housing was 13.1 m².¹⁷ Furthermore, according to a survey in 2009, Hong Kong had the lowest average residential floor space per person among 14 countries worldwide.³⁹ When compared to Australia, Hong Kong has only one-fifth of the average residential floor space per person. For the average new home size built in 2009, Hong Kong again had the smallest area,⁴⁰ which was less than one-fourth of those in Australia, Canada and the US. Our findings suggested that the small living space in Hong Kong is associated with a longer eye and a more minus refractive error. We speculate that the small home size and dense population may be two additional factors which are associated with the high prevalence of myopia in other East-Asian countries⁴¹ apart from other known factors.

This study was strong in several aspects. The participation rate (95%) was high because this research project was also a community service project, which did not further filter subjects within the sampled groups. The sampling method was modified to recruit a proportional number of subjects from districts of different population densities, so that the sample would reflect the characteristics of the population. We set out to make the questionnaire as simple and straightforward as possible so that parents could easily provide valid data. Qualified optometric personnel conducted all measurements in the study to ensure the accuracy of the results.

Yet, our study was not without limitations. A cycloplegic agent was not instilled because the data were collected on normal school days, and we did not want to interrupt the students' daily study. This may affect the accuracy of the auto-refraction as the subjects may accommodate, resulting in a more minus SER.¹⁹ However, the SER results were strongly correlated with the AL measurements (SER vs. AL: $r = -0.74$, $p < 0.001$), and hence could still identify the risk factors in the regression model. In addition, the data collection process adopted a self-reported questionnaire instead of an interview, which may hinder the data reliability to some extent. We tried to maximise the readability and ensure parents could understand the questionnaire without further explanation by inviting 10 laymen to answer the questionnaire. Our cross-sectional study could only establish the association between ocular parameters and living environment at a single time point. Further longitudinal studies shall be conducted to investigate the relationship between constricted living space and refractive error development.

Conclusion

In conclusion, there was an association between childhood refractive error and living environment, in terms of the size of home and the population density of the residential area. We speculate small homes and densely populated residential areas may be new types of 'visual pollutants' that associate with the high prevalence of myopia.

Disclosure

The authors report no conflicts of interest.

Acknowledgements

This study was supported by General Research Fund from the Research Grants Council of the Hong Kong Special Administrative Region, China (PolyU 5605/13M) and Internal Research Grants, The Hong Kong Polytechnic University (G-YBBS, Z0GF). The authors would like to

acknowledge Dr. J Neuville for proofreading the manuscript, and Dr. JX Lian for providing statistical advice.

References

1. Verkicharla PK, Ohno-Matsui K & Saw SM. Current and predicted demographics of high myopia and an update of its associated pathological changes. *Ophthalmic Physiol Opt* 2015; 35: 465–475.
2. Vu HT, Keeffe JE, McCarty CA & Taylor HR. Impact of unilateral and bilateral vision loss on quality of life. *Br J Ophthalmol* 2005; 89: 360–363.
3. Pan CW, Ramamurthy D & Saw SM. Worldwide prevalence and risk factors for myopia. *Ophthalmic Physiol Opt* 2012; 32: 3–16.
4. Fan DS, Lam DS, Lam RF *et al.* Prevalence, incidence, and progression of myopia of school children in Hong Kong. *Invest Ophthalmol Vis Sci* 2004; 45: 1071–1075.
5. Lam CS, Lam CH, Cheng SC & Chan LY. Prevalence of myopia among Hong Kong Chinese schoolchildren: changes over two decades. *Ophthalmic Physiol Opt* 2012; 32: 17–24.
6. Ip JM, Huynh SC, Robaei D *et al.* Ethnic differences in refraction and ocular biometry in a population-based sample of 11–15-year-old Australian children. *Eye (Lond)* 2008; 22: 649–656.
7. Twelker JD, Mitchell GL, Messer DH *et al.* Children's ocular components and age, gender, and ethnicity. *Optom Vis Sci* 2009; 86: 918–935.
8. Jeynes W. What we should and should not learn from the Japanese and other East Asian education systems. *Educ Policy* 2008; 22: 900–927.
9. He M, Zheng Y & Xiang F. Prevalence of myopia in urban and rural children in mainland China. *Optom Vis Sci* 2009; 86: 40–44.
10. Uzma N, Kumar BS, Khaja Mohinuddin Salar BM, Zafar MA & Reddy VD. A comparative clinical survey of the prevalence of refractive errors and eye diseases in urban and rural school children. *Can J Ophthalmol* 2009; 44: 328–333.
11. Ip JM, Rose KA, Morgan IG, Burlutsky G & Mitchell P. Myopia and the urban environment: findings in a sample of 12-year-old Australian school children. *Invest Ophthalmol Vis Sci* 2008; 49: 3858–3863.
12. Mutti DO, Mitchell GL, Moeschberger ML, Jones LA & Zadnik K. Parental myopia, near work, school achievement, and children's refractive error. *Invest Ophthalmol Vis Sci* 2002; 43: 3633–3640.
13. Saw SM, Chua WH, Hong CY *et al.* Nearwork in early-onset myopia. *Invest Ophthalmol Vis Sci* 2002; 43: 332–339.
14. Ip JM, Saw SM, Rose KA *et al.* Role of near work in myopia: findings in a sample of Australian school children. *Invest Ophthalmol Vis Sci* 2008; 49: 2903–2910.
15. Wu X, Gao G, Jin J *et al.* Housing type and myopia: the mediating role of parental myopia. *BMC Ophthalmol* 2016; 16: 151.

16. Census and Statistics Department. *2011 Population Census Summary Results: The Government of The Hong Kong Special Administrative Region*; 2012. <http://www.census2011.gov.hk/pdf/summary-results.pdf>, accessed 15/11/2016.
17. Hong Kong Housing Authority. *Housing in Figures 2015: The Government of The Hong Kong Special Administrative Region*; 2015. <https://www.housingauthority.gov.hk/en/commom/pdf/about-us/publications-and-statistics/HIF.pdf>, accessed 15/11/2016.
18. Society for Community Organization. *Research Report on Cage Homes, Cubicles, and Sub-divided flats: Hong Kong Society for Community Organization*; 2013. http://www.soco.org.hk/publication/private_housing/cagehome%20research%202013.doc, accessed 15/11/2016.
19. Fotedar R, Rochtchina E, Morgan I, Wang JJ, Mitchell P & Rose KA. Necessity of cycloplegia for assessing refractive error in 12-year-old children: a population-based study. *Am J Ophthalmol* 2007; 144: 307–309.
20. Tang WC, Tang YY & Lam CS. How representative is the 'Representative Value' of refraction provided by the Shin-Nippon NVision-K 5001 autorefractor? *Ophthalmic Physiol Opt* 2014; 34: 89–93.
21. Rubin DB. *Multiple Imputation for Nonresponse in Surveys*. John Wiley & Sons: New York, 2004.
22. Negrel AD, Maul E, Pokharel GP, Zhao J & Ellwein LB. Refractive Error Study in Children: sampling and measurement methods for a multi-country survey. *Am J Ophthalmol* 2000; 129: 421–426.
23. Maul E, Barroso S, Munoz SR, Sperduto RD & Ellwein LB. Refractive Error Study in Children: results from La Florida, Chile. *Am J Ophthalmol* 2000; 129: 445–454.
24. Pokharel GP, Negrel AD, Munoz SR & Ellwein LB. Refractive Error Study in Children: results from Mechi Zone, Nepal. *Am J Ophthalmol* 2000; 129: 436–444.
25. Murthy GV, Gupta SK, Ellwein LB *et al.* Refractive error in children in an urban population in New Delhi. *Invest Ophthalmol Vis Sci* 2002; 43: 623–631.
26. Naidoo KS, Raghunandan A, Mashige KP *et al.* Refractive error and visual impairment in African children in South Africa. *Invest Ophthalmol Vis Sci* 2003; 44: 3764–3770.
27. He M, Zeng J, Liu Y, Xu J, Pokharel GP & Ellwein LB. Refractive error and visual impairment in urban children in southern China. *Invest Ophthalmol Vis Sci* 2004; 45: 793–799.
28. He M, Huang W, Zheng Y, Huang L & Ellwein LB. Refractive error and visual impairment in school children in rural southern China. *Ophthalmology* 2007; 114: 374–382.
29. Ojaimi E, Rose KA, Smith W, Morgan IG, Martin FJ & Mitchell P. Methods for a population-based study of myopia and other eye conditions in school children: the Sydney Myopia Study. *Ophthalmic Epidemiol* 2005; 12: 59–69.
30. Rose KA, Morgan IG, Ip J *et al.* Outdoor activity reduces the prevalence of myopia in children. *Ophthalmology* 2008; 115: 1279–1285.
31. Smith EL, Kee CS, Ramamirtham R, Qiao-Grider Y & Hung LF. Peripheral vision can influence eye growth and refractive development in infant monkeys. *Invest Ophthalmol Vis Sci* 2005; 46: 3965–3972.
32. Smith EL, Hung LF & Huang J. Relative peripheral hyperopic defocus alters central refractive development in infant monkeys. *Vision Res* 2009; 49: 2386–2392.
33. Wallman J, Gottlieb MD, Rajaram V & Fugate-Wentzek LA. Local retinal regions control local eye growth and myopia. *Science* 1987; 237: 73–77.
34. Diether S & Schaeffel F. Local changes in eye growth induced by imposed local refractive error despite active accommodation. *Vision Res* 1997; 37: 659–668.
35. Tse DY, Lam CS, Guggenheim JA *et al.* Simultaneous defocus integration during refractive development. *Invest Ophthalmol Vis Sci* 2007; 48: 5352–5359.
36. Flitcroft DI. The complex interactions of retinal, optical and environmental factors in myopia aetiology. *Prog Retin Eye Res* 2012; 31: 622–660.
37. Saw SM, Nieto FJ, Katz J, Schein OD, Levy B & Chew SJ. Factors related to the progression of myopia in Singaporean children. *Optom Vis Sci* 2000; 77: 549–554.
38. Saw SM, Nieto FJ, Katz J, Schein OD, Levy B & Chew SJ. Familial clustering and myopia progression in Singapore school children. *Ophthalmic Epidemiol* 2001; 8: 227–236.
39. Lindsay W. *England: Shrink that footprint*; 2009. http://shrinkthatfootprint.com/wp-content/uploads/2013/04/Perca_pita.gif, accessed 15/11/2016.
40. Lindsay W. *England: Shrink that footprint*; 2009. <http://shrinkthatfootprint.com/wp-content/uploads/2013/04/Houseizem21.gif>, accessed 15/11/2016.
41. Ho A. *The unlivable dwellings in Hong Kong and the minimum living space*; 2015. <https://www.hongkongfp.com/2015/07/27/the-unlivable-dwellings-in-hong-kong-and-the-mini-mum-living-space/>, accessed 15/11/2016.